

**A MOMENT INEQUALITY
FOR THE MAXIMUM PARTIAL SUMS
WITH A GENERALIZED SUPERADDITIVE STRUCTURE**

Tibor Tómacs (EKTF, Hungary)

Abstract: F. A. Móricz, R. J. Serfling and W. F. Stout (1982) proved a moment inequality with superadditive function. The theorem of this paper extends this result to multidimensional sequence.

1. Notations

In the following \mathbb{Z} , \mathbb{N} , \mathbb{R} and d denotes the set of integers, positive integers, real numbers and a fixed positive integer. We define $\underline{1} = (1, 1, \dots, 1) \in \mathbb{N}^d$ and if $\underline{k} = (k_1, k_2, \dots, k_d) \in \mathbb{Z}^d$, $\underline{l} = (l_1, l_2, \dots, l_d) \in \mathbb{Z}^d$, $k_i \leq l_i$ for each $1 \leq i \leq d$ then $\underline{k} \leq \underline{l}$. The $\underline{k} < \underline{l}$ relation is defined similarly. If there exists an i index such that $k_i \geq l_i$ then we write $\underline{k} \not\leq \underline{l}$. Denote $|\underline{k}| = \prod_{i=1}^d k_i$ and let $\{X_{\underline{k}} : \underline{k} \in \mathbb{N}^d\}$ be a d -multiple sequence of random variables. $S_{\underline{n}}$ will denote the sum $\sum_{\underline{k} \leq \underline{n}} X_{\underline{k}}$ if $\underline{n} \in \mathbb{N}^d$, otherwise $S_{\underline{n}} = 0$. Finally $\mathbb{E}X$ will denote the expectation of the random variable X .

2. Preliminary results

Let $g: \mathbb{N}^2 \rightarrow \mathbb{R}$ be a nonnegative function. If $g(i, j) + g(j + 1, k) \leq g(i, k)$ for all $1 \leq i \leq j < k$ then we say that g is superadditive. F. A. Móricz, R. J. Serfling and W. F. Stout (1982) proved the next theorem: If $\{X_l : l \in \mathbb{N}\}$ sequence of random variables, $\alpha > 1$, $r \geq 1$, g is a superadditive function and

$$\mathbb{E} \left| \sum_{l=i}^j X_l \right|^r \leq g^\alpha(i, j)$$

for all $1 \leq i \leq j$ integers then there exists a constant $A_{\alpha, r}$ (what depends on α and r) such that for each $n \in \mathbb{N}$

$$\mathbb{E} \left(\max_{k \leq n} \left| \sum_{l=1}^k X_l \right| \right)^r \leq A_{\alpha, r} g^\alpha(1, n).$$

F. A. Móricz (1983) generalized the definition of superadditive function for d -dimension as follows. Let $g: \mathbb{N}^d \times \mathbb{N}^d \rightarrow \mathbb{R}$ be a nonnegative function. If

$$g(\underline{i}, \underline{j}) + g(\hat{\underline{i}}, \hat{\underline{j}}) \leq g(\underline{i}, \underline{j}), \quad (2.1)$$

where $\underline{i}, \underline{j} \in \mathbb{N}^d$, $\underline{i} \leq \underline{j}$, $1 \leq l \leq d$, $i_l \leq k_l \leq j_l$ and

$$\begin{aligned} \hat{\underline{i}} &= (i_1, \dots, i_{l-1}, k_l + 1, i_{l+1}, \dots, i_d), \\ \hat{\underline{j}} &= (j_1, \dots, j_{l-1}, k_l, j_{l+1}, \dots, j_d) \end{aligned}$$

then we say that the g is superadditive. F. A. Móricz (1983) proved the next theorem what is generalization of the previous theorem. If g is a superadditive function, $\alpha > 1$, $r \geq 1$ and

$$\mathbb{E} \left| \sum_{\underline{i} < \underline{l} \leq \underline{j}} X_{\underline{l}} \right|^r \leq g^\alpha(\underline{i}, \underline{j}) \quad (2.2)$$

for all $\underline{i}, \underline{j} \in \mathbb{N}^d$, $\underline{i} \leq \underline{j}$ then there exists a constant $A_{\alpha, r, d}$ (what depends on α, r and d) such that

$$\mathbb{E} \left(\max_{\underline{k} \leq \underline{n}} |S_{\underline{k}}| \right)^r \leq A_{\alpha, r, d} g^\alpha(\underline{1}, \underline{n})$$

for all $\underline{n} \in \mathbb{N}^d$. This paper discuss another generalization.

2. Main result

Theorem. Let $g: \mathbb{N}^d \times \mathbb{N}^d \rightarrow \mathbb{R}$ be a nonnegative function, $\alpha > 1$ and $r \geq 1$. Assume that for each $\underline{1} \leq \underline{i} \leq \underline{j} < \underline{k}$

$$g(\underline{i}, \underline{j}) + g(\underline{j} + \underline{1}, \underline{k}) \leq g(\underline{i}, \underline{k}), \quad (3.1)$$

and

$$\mathbb{E} |S_{\underline{j}} - S_{\underline{i}-\underline{1}}|^r \leq g^\alpha(\underline{i}, \underline{j}). \quad (3.2)$$

Then

$$\mathbb{E} \left(\max_{\underline{k} \leq \underline{n}} |S_{\underline{k}}| \right)^r \leq A_{\alpha, r} g^\alpha(\underline{1}, \underline{n}) \quad (3.3)$$

for all $\underline{n} \in \mathbb{N}^d$ where $A_{\alpha, r} = \left(1 - \frac{1}{2^{(\alpha-1)/r}}\right)^{-r}$.

Remark. This theorem is generalization of result of F. A. Móricz, R. J. Serfling and W. F. Stout (1982). We remark that condition (2.1) implies (3.1) on the other hand

if $X_{\underline{k}}$ ($\underline{k} \in \mathbb{N}^d$) nonnegative random variables then (3.2) implies (2.2) moreover the constant is not depending on d .

Proof of Theorem. Assume that $\underline{1} < \underline{N} = (N, N, \dots, N) \in \mathbb{N}^d$ and $\underline{n} \in \mathbb{N}^d$ where $\underline{n} \leq \underline{N}$ and $\underline{n} \not\leq \underline{N}$. If $|j| = 0$ then let $g(\underline{1}, \underline{j}) = 0$. With these notations, since

$$g(\underline{i}, \underline{j}) \leq g(\underline{i}, \underline{j}) + g(\underline{j} + \underline{1}, \underline{k}) \leq g(\underline{i}, \underline{k}) \quad \forall \underline{1} \leq \underline{i} \leq \underline{j} < \underline{k}, \quad (3.4)$$

there exists $m \geq 0$ integer having the property that

$$g(\underline{1}, \underline{m} - \underline{1}) \leq \frac{1}{2}g(\underline{1}, \underline{n}) \leq g(\underline{1}, \underline{m}), \quad (3.5)$$

where $\underline{m} = \underline{n} - m \cdot \underline{1}$. So if $\underline{m} < \underline{n}$ then

$$\frac{1}{2}g(\underline{1}, \underline{n}) + g(\underline{m} + \underline{1}, \underline{n}) \leq g(\underline{1}, \underline{m}) + g(\underline{m} + \underline{1}, \underline{n}) \leq g(\underline{1}, \underline{n}).$$

Consequently we have

$$g(\underline{m} + \underline{1}, \underline{n}) \leq \frac{1}{2}g(\underline{1}, \underline{n}), \quad \text{if } \underline{m} < \underline{n}. \quad (3.6)$$

Let us define sets

$$\begin{aligned} B &= \{\underline{k} \in \mathbb{N}^d : \underline{k} < \underline{m}\} \\ C &= \{\underline{k} \in \mathbb{N}^d : \underline{k} \leq \underline{n}, \underline{k} \not\leq \underline{m}, \underline{m} \not\leq \underline{k}\} \\ D &= \{\underline{k} \in \mathbb{N}^d : \underline{m} < \underline{k} \leq \underline{n}\} \end{aligned}$$

Let $\underline{k}_1 \in D$ such that $|S_{\underline{k}_1}| = \max_{\underline{k} \in D} |S_{\underline{k}}|$. If $D = \emptyset$ (other words $\underline{m} = \underline{n}$) then let $\underline{k}_1 = \underline{m}$. Let $\underline{k}_2 \in C$ such that $|S_{\underline{k}_2}| = \max_{\underline{k} \in C} |S_{\underline{k}}|$. With these notations we have

$$\begin{aligned} \max_{\underline{k} \leq \underline{n}} |S_{\underline{k}}| &= \max\{\max_{\underline{k} < \underline{m}} |S_{\underline{k}}|, |S_{\underline{k}_1}|, |S_{\underline{k}_2}|\} \leq \\ &= \max\{\max_{\underline{k} < \underline{m}} |S_{\underline{k}}|, |S_{\underline{m}}| + |S_{\underline{k}_1} - S_{\underline{m}}|, |S_{\underline{k}_2}|\} \leq \\ &= |S_{\underline{k}_2}| + \max\{\max_{\underline{k} < \underline{m}} |S_{\underline{k}}|, |S_{\underline{k}_1} - S_{\underline{m}}|\} \leq \\ &= |S_{\underline{k}_2}| + \left(\max_{\underline{k} < \underline{m}} |S_{\underline{k}}|^r + |S_{\underline{k}_1} - S_{\underline{m}}|^r \right)^{1/r}. \end{aligned} \quad (3.7)$$

The Minkowski's inequality states that

$$(\mathbb{E}|X + Y|^r)^{1/r} \leq (\mathbb{E}|X|^r)^{1/r} + (\mathbb{E}|Y|^r)^{1/r}$$

where X, Y random variables and $r \geq 1$. Therefore with $X = |S_{\underline{k}_2}|$ and $Y = \max_{\underline{k} < \underline{n}} |S_{\underline{k}}| - |S_{\underline{k}_2}|$ substitutions the Minkowski's inequality and (3.7) imply

$$\begin{aligned} \mathbb{E} \left(\max_{\underline{k} < \underline{n}} |S_{\underline{k}}|^r \right)^{1/r} &\leq (\mathbb{E} |S_{\underline{k}_2}|^r)^{1/r} + \left(\mathbb{E} \left| \max_{\underline{k} < \underline{n}} |S_{\underline{k}}| - |S_{\underline{k}_2}| \right|^r \right)^{1/r} \leq \\ &(\mathbb{E} |S_{\underline{k}_2}|^r)^{1/r} + \left(\mathbb{E} (\max_{\underline{k} < \underline{m}} |S_{\underline{k}}|^r) + \mathbb{E} |S_{\underline{k}_1} - S_{\underline{m}}|^r \right)^{1/r}. \end{aligned} \quad (3.8)$$

By condition (3.2) and (3.4) we get

$$(\mathbb{E} |S_{\underline{k}_2}|^r)^{1/r} \leq g^{\alpha/r}(\underline{1}, \underline{k}_2) \leq g^{\alpha/r}(\underline{1}, \underline{n}). \quad (3.9)$$

An elementary computation shows that $A_{\alpha, r} \geq 1$ so (3.2), (3.4) and (3.6) imply

$$\begin{aligned} \mathbb{E} |S_{\underline{k}_1} - S_{\underline{m}}|^r &\leq g^\alpha(\underline{m} + \underline{1}, \underline{k}_1) \leq g^\alpha(\underline{m} + \underline{1}, \underline{n}) \leq \\ &\frac{1}{2^\alpha} g^\alpha(\underline{1}, \underline{n}) \leq A_{\alpha, r} \frac{1}{2^\alpha} g^\alpha(\underline{1}, \underline{n}), \end{aligned} \quad (3.10)$$

if $D \neq \emptyset$ (what means $\underline{m} < \underline{k}_1$).

After these we prove the theorem by d -dimensional induction.

$$\mathbb{E} (\max_{\underline{k} \leq \underline{1}} |S_{\underline{k}}|^r) = \mathbb{E} |S_{\underline{1}}|^r \leq g^\alpha(\underline{1}, \underline{1}) \leq A_{\alpha, r} g^\alpha(\underline{1}, \underline{1})$$

therefore $\underline{n} = \underline{1}$ satisfies (3.3). Now, assume that (3.3) is true if $\underline{n} < \underline{N}$. Thus (3.5) implies

$$\mathbb{E} (\max_{\underline{k} < \underline{m}} |S_{\underline{k}}|^r) \leq A_{\alpha, r} g^\alpha(\underline{1}, \underline{m} - \underline{1}) \leq A_{\alpha, r} \frac{1}{2^\alpha} g^\alpha(\underline{1}, \underline{n}). \quad (3.11)$$

Finally by (3.8), (3.9), (3.10) and (3.11) we obtain

$$\mathbb{E} (\max_{\underline{k} \leq \underline{n}} |S_{\underline{k}}|^r)^{1/r} \leq g^{\alpha/r}(\underline{1}, \underline{n}) + \left(A_{\alpha, r} \frac{1}{2^{\alpha-1}} g^\alpha(\underline{1}, \underline{n}) \right)^{1/r} = A_{\alpha, r}^{1/r} g^{\alpha/r}(\underline{1}, \underline{n})$$

therefore (3.3) is true for each \underline{n} with $\underline{n} \leq \underline{N}$ and $\underline{n} \not\leq \underline{N}$. This completes the proof of the theorem.

References

- [1] F. A. MÓRICZ, R. J. SERFLING, W. F. STOUT, Moment and probability bounds with quasi-superadditive structure for the maximum partial sum, *The Annals of Probability* **Vol. 10, No.4** (1982), 1032–1040.

- [2] F. A. MÓRICZ, A general moment inequality for the maximum of the rectangular partial sums of multiple series, *Acta Math. Hung.*, **41** (3–4) (1983), 337–346.

Tibor Tómacs

Institute of Mathematics and Informatics
Károly Eszterházy Teachers' Training College
Leányka str. 4–6.
H-3300 Eger, Hungary